

# Mechanical ventilation & pulmonary complications in stroke

---

---

1. synchronized intermittent mandatory ventilation (SIMV) volume controlled ventilation (VCV)가 (pO<sub>2</sub> ~ 100 mmHg) (pCO<sub>2</sub> = 35 ~ 45 mmHg) pCO<sub>2</sub> 35 mmHg (jugular venous return) 60 mmHg 40 (hemispheric) bulbar dysfunction 가 12 cmH<sub>2</sub>O 가 90%, 10% [1], 6 ~ 30% 가 Succinylcholine 가 propofol, thiopental, etomidate 가 Midazolam 5 mg rocuronium 가 meta-analysis 가 [2].

2. bulbar dysfunction pressure support mode가

3.

4.

44% ,  
 2 ~ 3 가 .  
 가 , NIH  
 stroke 가 [3].  
 , vertebrobasilar  
 , [4].  
 (gastrostomy)  
 가 .  
 5.  
 30  
 58% (46 ~ 75%) 1/3  
 가 가

가  
 [5].  
 fatal midbrain herniation [1].

## REFERENCES

1. Berrouschot J. et al, Mechanical ventilation in patients with hemispheric ischemic stroke. Crit Care Med 2000;28:2956-2961.
2. Coplin WM. et al, Implications of extubation delay in brain-injured patients meeting standard weaning criteria. Am J Respir Crit Care Med 2000;161:1530-36.
3. Dziewas R. et al, Pneumonia in acute stroke patients fed by nasogastric tube. J Neurol Neurosurg Psychiatry 2004;75:852-856.
4. Hilker R. et al, Nosocomial pneumonia after acute stroke. Stroke 2003;34:975-981.
5. Holloway RG. et al, Prognosis and decision making in severe stroke. JAMA 2005;294:725-733.